

The Anomalous Chromomagnetic Dipole Moment of Top Quark

R. Martínez¹, J.-Alexis Rodríguez^{1,2} and M. Vargas¹

¹*Depto. de Física, Universidad Nacional, Bogotá, Colombia*

²*Centro Internacional de Física, Bogotá, Colombia*

()

We calculate the one loop corrections to the chromomagnetic dipole moment of the top quark in the framework of the Standard Model, the two Higgs doublet model and the Minimal Supersymmetric Standard Model. In the Standard Model we consider the QCD and the electroweak corrections generated by gluon and Higgs boson exchanges, respectively. In the Minimal Supersymmetric Standard Model, we calculate the SUSY- QCD corrections, including the mixing effects between the stop coming from the left- and right-handed sectors. We obtain that the Standard Model contribution is of the order of $-0.004 \leq \delta\kappa_g^t \leq -0.001$, and the values for diferents scenarios are of the order of $-0.001 \leq \delta\kappa_g^t \leq -0.01$.

Up to now, the Standard Model (SM) gives an adequate description of the data that involve the strong and electroweak interactions [1] . In particular, the discovery of the top quark at Fermilab by CDF and D0 [2], with a mass of 174 ± 6 GeV within the range allowed by the high precision tests of the SM [3], has showed that the SM is a good scheme that describes physics at these energies. However, the SM is not considered the final theory, and physicists think in the existence of new physics. In particular, since the fermion mass generation may be closely related to the electroweak symmetry breaking, one expects that some residual effects of the breaking mechanism to appear in accordance with the generated mass [4].

The top quark as the heaviest fermion in the model with a mass of the order of the symmetry breaking scale, opens a window to new physics, and one natural scenario is the modification of its interactions with the known gauge bosons. Future experimental runs will increase the number of produced $t\bar{t}$ pairs, allowing for the comparison of measured

differential cross sections with theoretical predictions of the many different models. In particular, one hopes to obtain detailed information on the couplings of the top quark [5].

Some $SU(2)_L \otimes U(1)_Y$ invariant operators, which represent new physics effects, give rise to couplings of the dipole moment type [5] [6]. In the case of strong interactions, there are some CP-invariant operators that contribute to the anomalous chromomagnetic dipole moment of the top quark, $\delta\kappa_g^t$. Although there are stringent experimental bounds on anomalous contributions to dipole moments of light fermions, it is not unreasonable to expect large anomalous moments of the heavy top quark [7].

The chromomagnetic dipole moment deserves special interest because this coupling is crucial in the production of top quark pairs by the annihilation processes $e^+e^- \rightarrow \bar{t}t$ and $\bar{q}q \rightarrow \bar{t}t$ [8]. For instance, the top quark anomalous couplings can modify the energy distribution of the extra gluon jet associated with top pair productions, and the production of top quark pairs in association with a hard gluon in future e^+e^- experiments provides a good opportunity to probe the anomalous ttg couplings [9].

This possibility has been considered, exploring the implications of scenarios like the SM [10], the two-Higgs-doublet model (2HDM) [11] and the different contributions of the Minimal Supersymmetric Standard Model (MSSM) [12]. More recently, indirect bounds for the anomalous vertex have been obtained from known experiments, such as CLEO for B decays, which was used to bound the magnetic and chromomagnetic dipole moment of the top quark within the context of a decoupled effective Lagrangian [13].

To naively estimate the contribution of new physics effects to $\delta\kappa_g^t$, we can remove the gluon line from the vertex diagram, and then get the mass term for the top quark. By dimensional analysis this can be written as

$$\delta\kappa_g^t \approx \frac{m_t^2}{\Lambda^2} \ , \quad (1)$$

where Λ is the scale of new physics. If new physics appears at $\Lambda = 1$ TeV, then $\delta\kappa_g^t \approx 0.031$. In the literature some models that give non-zero $\delta\kappa_g^t$ value have been considered. Technicolor models [17] predict

$$\delta\kappa_g^t \approx \frac{2m_t^2}{m_\omega^2} , \quad (2)$$

where m_ω is the mass of the techniscalar exchanged in the loop. For $m_\omega \geq 0.5$ TeV, $\delta\kappa_g^t \leq 0.25$ [10].

Our goal in this paper is to calculate explicitly the anomalous chromomagnetic dipole moment of the top quark in the framework of the SM, the 2HDM, and supersymmetric QCD.

The first point that we wish to make in this paper is the status of the SM with respect to the ttg anomalous chromomagnetic dipole moment which is defined as

$$\mathcal{L} = i\delta\kappa_g^t \frac{g_s}{2m_t} \bar{t} T^a \sigma_{\mu\nu} q^\nu t G^{\mu,a} \quad (3)$$

where g_s and T_a are the usual $SU(3)_C$ coupling and operator. In the case of QCD, the anomalous factor comes from the calculation at one loop level of the vertex ttg by considering the exchange of gluons, and the result is

$$\delta\kappa_g^t = -\frac{\alpha_s(m_t)}{12\pi} . \quad (4)$$

We found that the top quark contribution into the loop is decoupling [18], and the dependence of $\delta\kappa_g^t$ with respect to m_t is determined by the running coupling constant.

On the other hand, the most important SM contribution to the anomalous chromomagnetic dipole coupling is due to the exchange of neutral Higgs boson and would-be Goldstone boson. The contribution of this sector can be expressed as:

$$\delta\kappa_g^t = \frac{-\sqrt{2}G_F m_t^2}{8\pi^2} \left[\int_0^1 dx \frac{x - x^3}{x^2 - a(m_h^2)x + 1} + \int_0^1 dx \frac{-x^3 + 2x^2 - x}{x^2 - a(m_Z^2)x + 1} \right] \quad (5)$$

with $a(m_i^2) = 2 - m_i^2/m_t^2$, m_h the Higgs mass and m_Z the Z mass. The above contribution is showed as a function of the Higgs mass in figure 1 (solid line). Also we display the sum of QCD and SM contributions (dashed line). The horizontal dashed line is the QCD contribution from eq.(4).

Our next task is to calculate the contribution of the 2HDM [14] to the anomalous chromomagnetic dipole moment of the top quark. This model preserves the $SU(2)_L \otimes U(1)_Y$

SM symmetry, and includes two scalar isodoublets Φ_i . The Yukawa couplings of the scalars and fermions are constrained by the requirement that there are no flavor changing neutral interactions at tree level [15]. There are two possible arrangements for the Yukawa couplings: either one may couple all of the fermions to a single Higgs doublet (model I), or one may couple Φ_1 to the right-handed down-type quarks, and Φ_2 to the right-handed up-type quarks. This is the so-called model II, which includes the scalar sector of the MSSM as a particular case. Here, we will consider the latter possibility. The Yukawa Lagrangian for the quark sector has the form

$$-L_Y = \sum_{i,j=1}^3 \mathcal{D}_{ij} \bar{q}_{i,L} \Phi_1 d_{j,R} + \sum_{i,j=1}^3 \mathcal{U}_{ij} \bar{q}_{i,L} \tilde{\Phi}_2 u_{j,R} \quad (6)$$

where $\bar{q}_{i,L}$ are the left-handed quark doublets, $\mathcal{U}$ and $\mathcal{D}$ are the up and down quark mass matrices, respectively.

In this case, the vacuum expectation values of the Higgs fields Φ_i can be chosen to be both real and positive:

$$\langle \Phi_1 \rangle = \begin{pmatrix} 0 \\ \nu_1/\sqrt{2} \end{pmatrix} \quad , \quad \langle \Phi_2 \rangle = \begin{pmatrix} 0 \\ \nu_2/\sqrt{2} \end{pmatrix} \quad (7)$$

By diagonalizing the Higgs mass matrices one finds the physical Higgs states. In the charged sector, there are a charged would-be Goldstone boson and a charged scalar Higgs,

$$\begin{aligned} G^\pm &= \phi_1^\pm \cos \beta + \phi_2^\pm \sin \beta \quad , \\ H^\pm &= \phi_2^\pm \cos \beta - \phi_1^\pm \sin \beta \end{aligned} \quad (8)$$

where the mixing angle β is given by $\tan \beta = \nu_2/\nu_1$. Similarly, the imaginary parts of the neutral components mix to form a neutral would-be Goldstone boson and a CP-odd scalar Higgs,

$$\begin{aligned} G^0 &= \Im \phi_1^0 \cos \beta + \Im \phi_2^0 \sin \beta \quad , \\ A^0 &= \Im \phi_2^0 \cos \beta - \Im \phi_1^0 \sin \beta \quad . \end{aligned} \quad (9)$$

Finally, the real parts of the components mix to form a pair of neutral CP-even scalars Higgs,

$$H^0 = \sqrt{2}((\Re\phi_1^0 - \nu_1) \cos \alpha + (\Re\phi_2^0 - \nu_2) \sin \alpha) , \quad (10)$$

$$h^0 = \sqrt{2}((\Re\phi_2^0 - \nu_2) \cos \alpha - (\Re\phi_1^0 - \nu_1) \sin \alpha)$$

where the mixing angle α depends on the Higgs potential parameters.

The quantity $\tan \beta$ plays a central role in the theory because the Yukawa couplings are often proportional to either $\tan \beta$ or $\cot \beta$,

Higgs	$\bar{u}u$	$\bar{d}d$
H^0	$m_u \sin \alpha / \sin \beta$	$m_d \cos \alpha / \cos \beta$
h^0	$m_u \cos \alpha / \sin \beta$	$-m_d \sin \alpha / \cos \beta$
A^0	$m_u \cot \beta \gamma_5$	$m_d \tan \beta \gamma_5$

In the above, we summarize the Yukawa couplings in model II (we have omitted the overall factors $-ig/2m_W$ for H^0, h^0 and $-g/2m_W$ for A^0). We may note that $0 \leq \beta \leq \pi/2$ since we take $\nu_1, \nu_2 > 0$. In Appendix A of ref. [16] it is shown that this implies $-\pi/2 \leq \alpha \leq 0$. In the same Appendix various choices for independent parameters of the Higgs sector are outlined. Following ref. [16], we adopt $\tan \beta$, m_{A^0} as our independent parameters, and m_{H^0} , m_{h^0} , α can be written as

$$\begin{aligned} m_{H^0, h^0}^2 &= \frac{1}{2} \left[m_{A^0}^2 + m_Z^2 \pm \sqrt{(m_{A^0}^2 + m_Z^2)^2 - 4m_Z^2 m_{A^0}^2 \cos^2 2\beta} \right] , \\ \cos 2\alpha &= -\cos 2\beta \left(\frac{m_{A^0}^2 - m_Z^2}{m_{H^0}^2 - m_{h^0}^2} \right) , \\ \sin 2\alpha &= -\sin 2\beta \left(\frac{m_{H^0}^2 + m_{h^0}^2}{m_{H^0}^2 - m_{h^0}^2} \right) . \end{aligned} \quad (11)$$

The contribution of the 2HDM to the anomalous chromomagnetic dipole moment of the quark top comes from the exchange of the neutral Higgs h_o , A_o , and H_o bosons and the would-be Goldstone boson. The complete expression is given by

$$\begin{aligned} \delta\kappa_g^t &= \frac{\sqrt{2}G_F m_t^2}{8\pi^2} \left[\left(\frac{\sin \alpha}{\sin \beta} \right)^2 \int_0^1 dx \frac{x - x^3}{x^2 - a(m_{H^0}^2)x + 1} - \left(\frac{\cos \alpha}{\sin \beta} \right)^2 \int_0^1 dx \frac{x - x^3}{x^2 - a(m_{h^0}^2)x + 1} \right. \\ &\quad \left. + \cot^2 \beta \int_0^1 dx \frac{-x^3 + 2x^2 - x}{x^2 - a(m_{A^0}^2)x + 1} + \int_0^1 dx \frac{-x^3 + 2x^2 - x}{x^2 - a(m_z^2)x + 1} \right] . \end{aligned} \quad (12)$$

In figure 2, we display the contribution in terms of the A^0 Higgs mass for a top mass of 175 GeV and for some values of $\tan\beta$. There, we have already summed the pure QCD correction.

Our last task is to estimate the anomalous chromomagnetic dipole moment in the context of the MSSM [19]. We only consider the QCD contribution in the MSSM by the supersymmetrization of the standard QCD contribution, i.e. the exchange of gluinos and squarks. The squark sector presents mixing between left- and right-handed sectors and this squark mixing is particularly relevant in the case of stop because it is proportional to the top quark mass.

If we neglect family mixings, the mixing relation is given by

$$\begin{aligned}\tilde{t}_a &= \sum_{b=ij} R_{ab} \tilde{q}_b \\ R &= \begin{pmatrix} \cos\theta_t & \sin\theta_t \\ -\sin\theta_t & \cos\theta_t \end{pmatrix}\end{aligned}\tag{13}$$

where the rotation matrix diagonalize the corresponding stop mass matrix [19]

$$M_t^2 = \begin{pmatrix} m_{\tilde{t}_L}^2 & \Upsilon m_t \\ \Upsilon m_t & m_{\tilde{t}_R}^2 \end{pmatrix},\tag{14}$$

with the expressions

$$\begin{aligned}m_{\tilde{t}_L}^2 &= \widetilde{M}_Q + m_Z^2 \cos 2\beta \left(\frac{1}{2} - \frac{2}{3} \sin^2 \theta_W \right) + m_t^2, \\ m_{\tilde{t}_R}^2 &= \widetilde{M}_U + m_Z^2 \cos 2\beta \left(\frac{2}{3} \sin^2 \theta_W \right) + m_t^2, \\ \Upsilon &= \mu \cot \beta + A_t \widetilde{M} \ ,\end{aligned}\tag{15}$$

where $\widetilde{M}_Q$, $\widetilde{M}_U$ are the soft supersymmetry breaking terms, μ is the coefficient of the mixing Higgs term in the superpotential and A_t is the trilinear soft SUSY-breaking parameter [19]. In this way, stop $\tilde{t}_1$ and $\tilde{t}_2$ become the mass eigenstates, with a mixing angle θ_t and mass eigenstates given by

$$\tan 2\theta_t = \frac{2\Upsilon m_t}{m_{\tilde{t}_L}^2 - m_{\tilde{t}_R}^2}$$

$$m_{\tilde{t}_{1,2}} = \frac{1}{2} \left[m_{\tilde{t}_L}^2 + m_{\tilde{t}_R}^2 \mp \sqrt{(m_{\tilde{t}_L}^2 - m_{\tilde{t}_R}^2)^2 + 4\Upsilon^2 m_t^2} \right] . \quad (16)$$

To evaluate the anomalous chromomagnetic dipole moment in this framework, we first consider the exchange of $\tilde{t}_1$. Our result can be written as

$$\delta\kappa_g^t = \frac{4\alpha_s}{3\pi} \int_0^1 dx \frac{\cos^2 \theta_t (x^2 - 3x^3/2) - \sin^2 \theta_t (x^3/2) + \hat{m}_g \sin \theta_t \cos \theta_t (x^2 - x)}{x^2 - (1 - \hat{m}_g^2 + \hat{m}_{\tilde{t}_1}^2)x + \hat{m}_{\tilde{t}_1}^2}$$

$$+ \frac{3\alpha_s}{2\pi} \int_0^1 dx \frac{\cos^2 \theta_t (x^2 - 3x^3/2) + \sin^2 \theta_t (x^3/2) - \hat{m}_g \sin \theta_t \cos \theta_t (x^2 - x)}{x^2 - (1 + \hat{m}_g^2 - \hat{m}_{\tilde{t}_1}^2)x + \hat{m}_g^2} \quad (17)$$

where $\hat{m}_i = m_i/m_t$. The expression for the exchange of $\tilde{t}_2$ is obtained by changing $\sin \theta_t$ by $\cos \theta_t$, and vice-versa. Figure 3 shows the anomalous chromomagnetic dipole moment as a function of $\tilde{M}$ with $\mu = -30$ GeV and $m_g = 200$ GeV, for some different values of $\tan \beta$. In figure 4 we plot the anomalous chromomagnetic dipole moment of the top quark as a function of μ with the fixed values $\tilde{M} = \tilde{M}_Q = \tilde{M}_U = 500$ GeV, $A_t = 0$, $m_g = 200$ GeV and $m_t = 175$ GeV, and for different values of $\tan \beta$ (as in the other figures). Finally, figure 5 shows $\delta\kappa_g^t$ as a function of the gluino mass m_g , for the same values of $\tan \beta$.

In conclusion, we have calculated the chromomagnetic dipole moment of the top quark in the framework of the SM, the 2HDM with a discrete symmetry (the so-called model II), as well as the SUSY-QCD corrections. In all cases, we have added the pure QCD correction which is proportional to α_s , like QED for the $g - 2$ factor. In the SM the main contribution comes from the neutral Higgs particle.

A naive estimation for the contribution of new physics gives $\delta\kappa_g^t \approx 0.031$ for $\Lambda = 1$ TeV, and $\delta\kappa_g^t \approx 0.25$ for a techniscalar mass of the order of 0.5 TeV [10]. Using the effective Lagrangian approach in the $b \rightarrow s\gamma$ process was found that $\delta\kappa_g^t \approx 0.066$ [13]. In this paper, we have found that the SM contribution is of the order of $-0.004 \leq \delta\kappa_g^t \leq -0.001$. This result is similar to the one obtained in ref. [21] for $\sqrt{s} = 500$ GeV. The values that we have found for different scenarios like 2HDM and SUSY-QCD are of the order of $-0.001 \leq \delta\kappa_g^t \leq -0.01$. In ref. [22] it was found that $ee \rightarrow \bar{t}t g$ processes at the Next Linear Collider (NLC)

will be sensitive to $-0.027 \leq \delta\kappa_g^t \leq 0.026$ which will allow to test physics beyond the SM through this dipole moment.

We acknowledge helpful discussions with M.A. Perez and F. Larios. The present work was complete during a visit to the Physics Department of CINVESTAV. It was supported by COLCIENCIAS and CONACyT. M. V. thanks Fundación Mazda para el Arte y la Ciencia for his fellowship.

- [1] S. Weinberg, Phys. Rev. Lett. **19**, 1264 (1967); A. Salam in Elementary Particle Theory ed. N. Southholm (Almquist and Wiksell, Stockholm (1968), pp 367; S. Glashow, Nucl. Phys. **22**, 579 (1961).
- [2] F. Abe et. al (CDF), Phys. Rev. Lett. **74**, 2626 (1995); S. Abachi *et. al.* (D0), Phys. Rev. Lett. **74**, 2632 (1995).
- [3] J. Bernabeu, A. Pich and A. Santamaria, Nucl. Phys. **B363**, 160 (1991)
- [4] R. D. Peccei, S. Peris and X. Zhang, Nucl. Phys. **B349**, 305 (1991); T. Han, R. D. Peccei and X. Zhang, Nucl. Phys. **B454**, 527 (1995); R. S. Chivucula *et. al.*, Phys. Lett. **B311**, 157 (1993); E. Malkawi and C.-P. Yuan, Phys. Rev. **D50**, 446 (1994); E. Malkawi and T. Tait, Phys. Rev. **D54**, 5758 (1996).
- [5] G. Kane, G. Ladinsky and C.-P. Yuan, Phys. Rev. **D45**, 1531 (1992); D. Carlson, E. Malkawi and C.-P. Yuan, Phys. Lett. **B337**, 145 (1994); P. D. Peccei and S. Peris, Phys. Rev. **D44**, 809 (1991); F. Larios and C.-P. Yuan, Phys. Rev. **D55**, 7218 (1997).
- [6] W. Buchmuller and D. Wyler, Nucl. Phys. **B268**, 621 (1986).
- [7] P. Haberl, O. Nachtmann and A. Wilch, Phys. Rev. **D53**, 4875 (1996).
- [8] D. Silverman, Phys. Rev. **D54**, 5562 (1996); T. G. Rizzo, Phys. Rev. **D53**, 2326 (1996).

- [9] T. Rizzo, hep-ph/9605361; hep-ph/9609311 and references cited there in.
- [10] D. Atwood, A. Kagan and T. Rizzo, Phys. Rev. **D52**, 6264 (1995); K. Cheung, Phys. Rev. **D53**, 3604 (1996)
- [11] A. Stange and S. Willenbrock, Phys. Rev. **D48**, 2054 (1993).
- [12] J. M. Yang and C. S. Li, Phys. Rev. **D54**, 4380 (1996); *ibid.*, **D52**, 1541 (1995); C. S. Li, H. Y. Zhou, Y. L. Zhu and J. M. Yang, hep-ph/9606271; S. Alam, K. Hawigara and S. Matsumoto, Phys. Rev. **D55**, 1307 (1997).
- [13] R. Martinez and J.-Alexis Rodriguez, Phys. Rev. **D55**, 3212 (1997); R. Martinez, M. A. Perez and J. J. Toscano, Phys. Lett. **B340**, 91 (1994).
- [14] S. Dawson, H. Haber, J. Gunion and G. Kane, Higgs Hunter's Guide, Frontiers in Physics, Addison Wesley, 1990; and references cited there in.
- [15] S. L. Glashow and S. Weinberg, Phys. Rev. **D15**, 1958 (1977); E. A. Paschos, Phys. Rev. **D15**, 1966 (1977).
- [16] J. F. Gunion and H. E. Haber, Nucl. Phys. **B278**, 449 (1986).
- [17] A. K. Kagan, Phys. Rev. **D51**, 6196 (1995).
- [18] T. Appelquist and J. Carazzone, Phys. Rev. **D11**, 2856 (1975).
- [19] H. Haber and G. Kane, Phys. Rep. **C117**, 75 (1985).
- [20] W. Boer *et. al.*, hep-ph/9609209; G. Altarelli, hep-ph/9611239.
- [21] J. Bernabéu *et. al.*, Phys. Rev. **D53**, 5222 (1996).
- [22] T. G. Rizzo, hep-ph/9605361.

Figure Captions

Figure 1. $\delta\kappa_g^t$ as a function of the Higgs mass. Horizontal dashed line is the QCD correction. The solid-line is the SM Higgs contribution, and dotted is the sum of QCD and Higgs SM contributions.

Figure 2. $\delta\kappa_g^t$ as a function of the Higgs mass in 2HDM for three different values of $\tan\beta = 1.5, 5$ and 15 corresponding to solid, dashed and dotted curves, respectively.

Figure 3. The same as the fig. 2 for the MSSM contribution as a function of $\tilde{M}$ with $\mu = -30$ GeV, $A_t = 0$ and $m_g = 200$ GeV.

Figure 4. The same as the fig. 3 as a function of μ instead, with $\tilde{M} = 500$ GeV, $A_t = 0$ and $m_g = 200$ GeV.

Figure 5. $\delta\kappa_g^t$ as a function of the gluino mass for $\mu = -30$ GeV, $A_t = 0$ and $\tilde{M} = 500$ GeV. $\tan\beta$ takes the same values of the fig. 2.

